

#1 1/2

Patent

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Jose Luis Castro Pineiro, et al.

Serial No.: To be assigned

Case No.: T1427

Filed: To be assigned

Art Unit:

Examiner:

For: Indole and indoline derivatives as 5-HT6 selective ligands



The Honorable Assistant Commissioner for Patents
Washington, D.C. 20231

SUBMITTAL OF PRIORITY DOCUMENT


Sir:

Enclosed please find a certified copy of the following priority document for
submittal with respect to the above-noted application:

Foreign/PCT Application - Great Britain 9820113.0 15/09/98.

Respectfully submitted,

By


James L. McGinnis
Reg. No. 34,387
Attorney for Applicant

MERCK & CO., Inc.
P.O. Box 2000
Rahway, New Jersey 07065-0907
(732) 594-0641

Date:

EXPRESS MAIL CERTIFICATE
DATE OF DEPOSIT Sept 9, 1999
EXPRESS MAIL NO. 9607116342105
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS
BEING DEPOSITED WITH THE UNITED STATES POSTAL
SERVICE AS EXPRESS MAIL "POST OFFICE TO
ADDRESSEE" BEFORE 5 P.M. ON THE ABOVE DATE IN
AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER
FOR PATENTS, WASHINGTON, D.C. 20231.
MAILED BY Althea Smith
DATE Sept. 9, 1999

THIS PAGE BLANK (USPTO)

1. THE
 2. THE
 3. THE
 4. THE
 5. THE
 6. THE
 7. THE
 8. THE
 9. THE
 10. THE
 11. THE
 12. THE
 13. THE
 14. THE
 15. THE
 16. THE
 17. THE
 18. THE
 19. THE
 20. THE
 21. THE
 22. THE
 23. THE
 24. THE
 25. THE
 26. THE
 27. THE
 28. THE
 29. THE
 30. THE
 31. THE
 32. THE
 33. THE
 34. THE
 35. THE
 36. THE
 37. THE
 38. THE
 39. THE
 40. THE
 41. THE
 42. THE
 43. THE
 44. THE
 45. THE
 46. THE
 47. THE
 48. THE
 49. THE
 50. THE
 51. THE
 52. THE
 53. THE
 54. THE
 55. THE
 56. THE
 57. THE
 58. THE
 59. THE
 60. THE
 61. THE
 62. THE
 63. THE
 64. THE
 65. THE
 66. THE
 67. THE
 68. THE
 69. THE
 70. THE
 71. THE
 72. THE
 73. THE
 74. THE
 75. THE
 76. THE
 77. THE
 78. THE
 79. THE
 80. THE
 81. THE
 82. THE
 83. THE
 84. THE
 85. THE
 86. THE
 87. THE
 88. THE
 89. THE
 90. THE
 91. THE
 92. THE
 93. THE
 94. THE
 95. THE
 96. THE
 97. THE
 98. THE
 99. THE
 100. THE



The
**Patent
Office**



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 19 August 1999

THIS PAGE BLANK (USPTO)


Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference	T1427PV		
2. Patent application number (The Patent Office will fill in this part)	<div style="text-align: right;"> <small>14 SEP 1998 E390474-1 D02432</small> <small>F01/7700 25.00 - 9820113.0</small> </div> <div style="text-align: center; font-size: 2em; font-weight: bold;">9820113.0</div>		
3. Full name, address and postcode of the or of each applicant (underline all surnames)	Merck Sharp & Dohme Limited Hertford Road, Hoddesdon Hertfordshire EN11 9BU United Kingdom		
Patents ADP number (if you know it)	00597799001 ✓		
If the applicant is a corporate body, give the country/state of its incorporation	United Kingdom		
4. Title of the invention	Therapeutic Agents		
5. Name of your agent (if you have one)	Dr. J. Thompson		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	Merck & Co., Inc. European Patent Department Terlings Park Eastwick Road Harlow Essex CM20 2QR		
Patents ADP number (if you know it)	4392742002 ✓		
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority Application number (if you know it)	Date of filing (day/month/year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application		Date of filing (day/month/year)
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:	Yes		
a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d))			



Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document


Continuation sheets of this form	0
Description	48
Claim(s)	0
Abstract	0
Drawing(s)	0



10. If you are also filing any of the following, state how many against each item.

Priority documents	-
Translations of priority documents	-
Statement of inventorship and right to grant of a patent (Patents Form 7/77)	-
Request for preliminary examination and search (Patents Form 9/77)	-
Request for substantive examination (Patents Form 10/77)	-
Any other documents (please specify)	Fee Sheet

11. I/We request the grant of a patent on the basis of this application.

Signature		Date	15 September 1998
Dr. J. Thompson			

12. Name and daytime telephone number of person to contact in the United Kingdom

Dr. J. Thompson	01279 440172
-----------------	--------------

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

THERAPEUTIC AGENTS

The present invention relates to classes of indole derivatives which bind selectively to 5-HT₆ receptors, and hence are potentially useful in the treatment of a number of clinical conditions believed to be sensitive to the selective agonism or antagonism of the 5-HT₆ receptors. Certain of the compounds are novel.


The actions of the neurotransmitter 5-hydroxytryptamine (5-HT) are mediated through a number of receptor families termed 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆, and 5-HT₇ (see D. Hoyer et al., *Pharmacol. Rev.*, 1994, 46, 157-204). Although the functions of the 5-HT₅, 5-HT₆ and 5-HT₇ receptors are less well understood than the functions of the other 5-HT receptors, it is generally accepted that compounds which selectively interfere with 5-HT-mediated signal transduction are important novel drug targets. In particular, 5-HT₆ selective ligands have been identified as potentially useful in the treatment of certain CNS disorders such as Parkinson's disease, Huntingdon's disease, anxiety, depression, manic depression, psychosis, epilepsy, obsessive compulsive disorders, migraine, Alzheimers disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia and bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Such compounds are also expected to be of use in the treatment of certain GI (gastrointestinal) disorders such as IBS (Irritable Bowel Syndrome). (See WO98/27058, WO98/27081, Drug News Perspectives, 1997, 10, 214 and A.J. Sleight et al., *Serotonin ID Research Alert*, 1997, 2(3), 115-8).

A wide variety of 5-HT analogues incorporating an indole nucleus have been prepared and tested for pharmacological effects, but hitherto none has been shown to be a selective ligand for 5-HT₆ receptors. WO-A-98/27058 and WO-A-98/27081 disclose, respectively, sulphonamides


and carboxamides which are 5-HT₆ receptor antagonists, but none of the compounds disclosed comprises an indole or indoline nucleus. Similarly, certain benzenesulphonamides of non-indole nature are disclosed as selective 5-HT₆ antagonists in EP-A-815861 and in *Br.J.Pharmacol.*, 1998, 124, 556-562.

Many analogues of 5-HT are known, comprising a 1-acyl- or 1-arylsulphonyl-3-(2-aminoethyl) indole nucleus (see, for example, US Patents Nos. 3,481,953 and 3,489,429). Fused-ring analogues, such as trans-4-dimethylamino-1-(4-methylbenzenesulphonyl)-1,3,4,5-tetrahydrobenz[cd] indol-5-ol, are also known (*J.Chem.Soc., Perkin Trans.I*, 1973, 438-42). Analogues comprising a 4-(2-N,N-dialkylaminoethyl)indole nucleus, unsubstituted in the 1-position, have been shown to be dopamine agonists (*Eur. J. Med. Chem.*, 1991, 26, 473-5; *J. Med. Chem.*, 1984, 27, 386-9). However, the 1-substituted derivatives disclosed herein are believed to be novel.


The present invention, in a first aspect, provides the use of a compound, or a pharmaceutically acceptable salt or prodrug thereof, for the manufacture of a pharmaceutical composition for the treatment and/or prevention of clinical conditions for which selective agonism or antagonism of 5-HT₆ receptors is indicated, said compound being an indole or indoline derivative having a structure in accordance with Formula I, II or III:



(I)



(II)



wherein

n is 1 or 2;

5 p is 0,1,2 or 3;

q is 0,1,2,3 or 4;

R¹ and R² independently represent hydrogen, C₁₋₆ alkyl or aryl (C₁₋₆)alkyl, or together represent the atoms necessary to complete a heterocycloalkyl group comprising the nitrogen atom to which R¹ and R² are attached;

R³ represents hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl(C₁₋₆)alkyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl or C₁₋₆ alkylcarbonyl;

15 R⁴ represents arylsulphonyl, heteroarylsulphonyl, C₁₋₆ alkylsulphonyl, di(C₁₋₆)alkylaminosulphonyl, arylcarbonyl, C₁₋₆ alkylcarbonyl heteroarylcarbonyl or C₁₋₆ alkoxycarbonyl;

each R⁵ independently represents hydroxy, C₁₋₆ alkoxy, aryl(C₁₋₆)alkoxy, nitrile or halogen;

R⁶ represents hydrogen, hydroxy or C₁₋₆ alkoxy; and

20 -A-B- represents -C=C- or -CH-CH-.

Also in its first aspect, the invention further provides a method for the treatment and/or prevention of clinical conditions for which selective agonism or antagonism of 5-HT₆ receptors is indicated comprising administering to a patient in need of such treatment an effective amount

of a compound in accordance with Formulae I, II or III, or a pharmaceutically acceptable salt or a prodrug thereof.

Compounds of Formula I wherein n,p, and R¹ - R⁵ are defined as before, and salts and prodrugs thereof, are novel and constitute a second
5 aspect of the invention.

In its second aspect, the invention also provides a pharmaceutical composition comprising one or more compounds of Formula I, or pharmaceutically acceptable salts or prodrugs thereof, in association with a pharmaceutically-acceptable carrier.

10 Also in its second aspect, the invention further provides a compound of Formula I or a pharmaceutically acceptable salt or a prodrug thereof for use in therapy.

The compounds of Formulae I-III are selective ligands for 5-HT₆ receptors, having a 5-HT₆ receptor (rat or human) binding affinity (K_i),
15 when measured in cell lines expressing cloned recombinant 5-HT₆ receptors, of less than 1μM, typically less than 100nM, and in preferred embodiments less than 10nM, and having a selective affinity for 5-HT₆ receptors relative to 5-HT₅ and/or 5-HT₇ receptors of at least 3-fold, typically at least 10-fold, and in preferred embodiments at least 100-fold.

20 In Formulae I-III, one or more substituents may be present on any alkyl or aryl group represented by any of R¹ - R⁵, or on any alkyl or aryl moiety of a group represented by any of R¹ - R⁵. Preferred substituents include C₁₋₆ alkyl, halogen, hydroxy and C₁₋₆ alkoxy.

As used herein, the expression "C₁₋₆ alkyl" includes methyl and
25 ethyl groups, and straight-chained, branched or cyclic propyl, butyl, pentyl and hexyl groups. Particular alkyl groups are methyl, ethyl, *n*-propyl, isopropyl and *tert*-butyl. Derived expressions such as "C₁₋₆ alkoxy", "C₁₋₆ alkylthio" and "C₁₋₆ alkylamino" are to be construed accordingly.

The expression "C₂₋₆ alkenyl" as used herein refers to straight-
30 chained and branched alkenyl groups containing from 2 to 6 carbon atoms. Typical examples include vinyl, allyl, dimethylallyl and butenyl groups.

The expression "C₂₋₆ alkynyl" as used herein refers to straight-chained and branched alkynyl groups containing from 2 to 6 carbon atoms. Typical examples include ethynyl and propargyl groups.

Typical aryl groups include phenyl and naphthyl.

5 The expression "aryl(C₁₋₆)alkyl" as used herein includes benzyl, phenylethyl, phenylpropyl and naphthylmethyl.

Suitable heterocycloalkyl groups include azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny groups.

Suitable heteroaryl groups include pyridiny, quinoliny,
10 isoquinoliny, pyridaziny, pyrimidiny, pyraziny, furyl, benzofuryl, dibenzofuryl, thienyl, benzthienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl and tetrazolyl groups.

The term "halogen" as used herein includes fluorine, chlorine,
15 bromine and iodine, especially chlorine or fluorine.

For use in medicine, the salts of the compounds of Formulae I-III will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds of Formulae I-III or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable
20 salts of the compounds of Formulae I-III include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic
25 acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

Furthermore, where the compounds of Formulae I-III carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable
30 organic ligands, e.g. quaternary ammonium salts.

The present invention includes within its scope the use of prodrugs of the compounds of Formulae I-III above. In general, such prodrugs will be functional derivatives of the compounds of Formulae I-III which are readily convertible *in vivo* into the required compounds of Formulae I-III.

5 Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in *Design of Prodrugs*, ed. H. Bundgaard, Elsevier, 1985.

Certain compounds according to the present invention may be capable of existing as tautomeric forms. It is to be understood that all

10 possible tautomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

Where the compounds according to the invention have at least one asymmetric centre, they may accordingly exist as enantiomers. Where the compounds according to the invention possess two or more asymmetric

15 centres, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

In Formulae I-III, suitable separate identities for R¹ and R² include hydrogen, methyl, ethyl, propyl and benzyl, and suitable identities for R¹

20 and R² in combination include pyrrolidinyl, piperidinyl, piperazinyl, 4-methylpiperazinyl and morpholinyl.

Suitable identities for R³ include hydrogen, methyl, ethyl, benzyl, allyl, propargyl, benzoyl, phenyl, thienyl and furoyl.

Suitable identities for R⁴ include benzenesulphonyl, naphthalene-2-

25 sulphonyl, o-, m- or p-toluenesulphonyl, o-, m- or p-chlorobenzenesulphonyl, o-, m- or p-methoxybenzenesulphonyl, methanesulphonyl, dimethylaminosulphonyl, thienylsulphonyl, benzoyl, acetyl, furoyl and *tert*-butoxycarbonyl.

Suitable identities for R⁵ include hydroxy, methoxy, ethoxy,

30 propoxy, benzyloxy, nitrile, fluorine, chlorine and bromine. Preferably, there is no more than one R⁵ substituent (i.e. p and q are 0 or 1), and when

a single R⁵ substituent is present, it is preferably in the para-position relative to the indole nitrogen.

In the compounds of Formula I, p is preferably zero; R¹ and R² are preferably identical and represent hydrogen or methyl; R³ preferably


represents hydrogen or benzoyl; and R⁴ preferably represents

arylsulphonyl or dimethylaminosulphonyl. Examples of specific

compounds in accordance with Formula I include:
2-[1-(benzenesulphonyl)-1*H*-indol-4-yl]ethylamine;
N,N-dimethyl 2-[1-(benzenesulphonyl)-1*H*-indol-4-yl]ethylamine;
N,N-dimethyl 2-[1-(dimethylamino)sulphonyl-1*H*-indol-4-yl]ethylamine;
N,N-dimethyl 3-[1-(benzenesulphonyl)-1*H*-indol-4-yl]propylamine; and
N,N-dimethyl 2-[1-(benzenesulphonyl)-2-benzoyl-1*H*-indol-4-yl]
ethylamine.

In the compounds of Formula II, preferably R¹ and R² are identical and represent hydrogen or methyl, or together complete a pyrrolidinyl, piperidinyl, piperazinyl or 4-methylpiperazinyl ring; R³ preferably represents hydrogen or methyl; R⁴ preferably represents arylsulphonyl, thienylsulphonyl, benzoyl or *tert*-butoxycarbonyl; R⁵ preferably represents hydroxy, methoxy, benzyloxy or nitrile; and q is zero or 1.

A sub-class of compounds in accordance with Formula II is defined by Formula II(a):



(II(a))

where R³, R⁴ and R⁵ have the same meanings as before.

Specific examples of compounds in accordance with Formula II(a) include:-

N,N-dimethyl 2-[1-(benzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine;

5 *N,N*-dimethyl 2-[5-methoxy-1-(4-methylbenzenesulphonyl)-1*H*-indol-3-yl]ethylamine;

N,N-dimethyl 2-[1-(4-chlorobenzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine;

10 *N,N*-dimethyl 2-[1-(3-chlorobenzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine;

N,N-dimethyl 2-[5-methoxy-1-(2-naphthalenesulphonyl)-1*H*-indol-3-yl]ethylamine;

N,N-dimethyl 2-[5-methoxy-1-(4-methoxybenzenesulphonyl)-1*H*-indol-3-yl]ethylamine;

15 *N,N*-dimethyl 2-[1-(2-chlorobenzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine;

N,N-dimethyl 2-(1-benzoyl-5-methoxy-1*H*-indol-3-yl)ethylamine;

N,N-dimethyl 2-[5-methoxy-1-(2-thiophenesulphonyl)-1*H*-indol-3-yl]ethylamine;

20 *N,N*-dimethyl 2-(1-benzenesulphonyl-5-methoxy-2-methyl-1*H*-indol-3-yl)ethylamine;

N,N-dimethyl 2-(1-benzenesulphonyl-1*H*-indol-3-yl)ethylamine;

N,N-dimethyl 2-(1-methylsulphonyl-1*H*-indol-3-yl)ethylamine;

N,N-dimethyl 2-(5-methoxy-1-methylsulphonyl-1*H*-indol-3-yl)ethylamine;


25 [3-(2-dimethylamino-ethyl)-5-hydroxy-1*H*-indol-1-yl]phenylmethanone;
3-(2-dimethylamino-ethyl)-5-hydroxy-1*H*-indole-1-carboxylic acid *tert*-butyl ester;

N,N-dimethyl 2-(1-benzenesulphonyl-5-benzyloxy-1*H*-indol-3-yl)ethylamine;

30 *N,N*-dimethyl 2-(1-benzenesulphonyl-5-hydroxy-1*H*-indol-3-yl)ethylamine;
and

N,N-dimethyl 2-(1-benzenesulphonyl-5-cyano-1*H*-indol-3-yl)ethylamine.

A further sub-class of compounds in accordance with Formula II is defined by Formula II(b):




(II(b))

where R^1 and R^2 have the same meanings as before, and Ar represents an aryl group.

Specific examples of compounds in accordance with Formula II(b) include:

2-[1-(benzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine;
 1-benzenesulphonyl-5-methoxy-3-[(2-pyrrolidin-1-yl)ethyl]-1*H*-indole;
 1-benzenesulphonyl-5-methoxy-3-[(2-piperidin-1-yl)ethyl]-1*H*-indole; and
 1-benzenesulphonyl-5-methoxy-3-[(2-piperazin-1-yl)ethyl]-1*H*-indole.

A third sub-class of compounds in accordance with Formula II is defined by Formula II(c):



(II(c))

where $R^1 - R^5$ have the same meanings as before.

Specific examples of compounds in accordance with Formula II(c) include:

- 5 *N,N*-dimethyl 2-(1-benzenesulphonyl-5-methoxy-2,3-dihydro-1*H*-indol-3-yl)ethylamine.

In the compounds of Formula III, R^1 and R^2 are preferably identical and represent hydrogen or methyl; R^3 preferably represents hydrogen; R^4 preferably represents arylsulphonyl, especially *p*-toluenesulphonyl; R^6 preferably represents hydroxy or methoxy; and *p* is preferably zero.

Specific examples of compounds in accordance with Formula III include: *trans*-4-dimethylamino-1-(4-methylbenzenesulphonyl)-1,3,4,5-tetrahydro-benz[*cd*]indol-5-ol; and 4-dimethylamino-5-methoxy-1-(4-methylbenzenesulphonyl)-1,3,4,5-tetrahydro-benz[*cd*]indole.

The invention also provides or uses pharmaceutical compositions comprising one or more compounds of Formulae I-III in association with a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with

a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing
5 a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage
10 forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. The tablets
15 or pills of the pharmaceutical composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer
20 which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.


25 The liquid forms in which the pharmaceutical compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and
30 similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as


tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

In the treatment of the relevant clinical conditions, such as depression, Alzheimer's disease, Parkinson's disease and schizophrenia, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.05 to 100 mg/kg per day, and especially about 0.05 to 5 mg/kg per day.

The compounds may be administered on a regimen of 1 to 4 times per day.


A representative approach to the preparation of compounds of Formula I in which $n=1$ is illustrated in the following reaction scheme, in which the starting material (methyl indole-4-carboxylate) is prepared as described in J. Org. Chem., 1980, 45, 3350.





In step (1), reaction of the indole derivative with R⁴Cl in the presence of base introduces the desired R⁴- substituent in the 1-position (where R⁴ has the same definition as before). Alternatively, and in particular when R⁴ represents *tert*-alkoxycarbonyl, this step may be carried out last. In step (2), the methyl ester is first reduced to the corresponding hydroxymethyl derivative, e.g. by means of diisobutyl aluminium hydride, then oxidised to the aldehyde IV using manganese (IV) oxide. In step (3), base-catalysed condensation of the aldehyde with nitromethane affords the 2-nitroethenyl derivative, which may be reduced (e.g. by zinc amalgam) to the 2-aminoethyl derivative (step (4)). If desired, mono- or dialkylation of the primary amino group may be carried out by standard methods. The preferred dimethylamino derivatives (V) are most conveniently prepared by reductive alkylation of the primary amino with formaldehyde in the presence of sodium cyanoborohydride (step (5)).


The above-described indole-4-carboxaldehyde derivatives (IV) are suitable starting materials for the preparation of compounds of Formula I in which n=2, as shown in the following reaction scheme; where R¹, R² and R⁴ have the same meanings as before:



In step (1), reaction between the aldehyde (IV) and (carbethoxymethylene) triphenyl phosphorane provides the ethyl ester of the acrylic acid derivative, which is subsequently reduced by catalytic hydrogenation over Pd/C to the corresponding propionic ethyl ester in step (2). In step (3), reaction of the ester with R^1R^2NH provides the corresponding amide, which is then reduced to the amine (VI) by lithium aluminium hydride (step (4)). Alternatively, the same sequence of reactions may be performed on the corresponding N-unsubstituted indoles, with attachment of the R^4 group via reaction with R^4-Cl as a final step. This is preferable when R^4 represents alkylcarbonyl, *tert*-alkoxycarbonyl, arylcarbonyl, or heteroarylcarbonyl.

Compounds of Formula I in which R^3 represents alkyl, aryl or arylalkyl may be prepared by the routes shown above, but starting from the appropriate 2-substituted analogue of methyl indole-4-carboxylate.

Compounds of Formula II in which R⁵ is other than OH may be prepared from the intermediates VII(a) and VII(b) as shown in the following reaction scheme:




where R¹, R², R³ and R⁴ have the same meanings as before, and R^{5a} represents any group represented by R⁵ except hydroxy.

In the event that either or both of R¹ and R² represents hydrogen, it is necessary to protect the exocyclic amino group as the t-butoxycarbonyl (Boc) derivative prior to functionalising the ring nitrogen, the Boc group being subsequently cleaved under acid conditions.

Compounds of Formula II in which R⁵ represents hydroxy may be prepared by hydrogenolysis of the corresponding compounds in which R⁵ represents benzyloxy.

The intermediates VII(b) in which -A-B- represents -CH-CH- may be prepared by catalytic hydrogenation of the intermediates VII(a) in which -A-B- represents -C=C-. Thus, the compounds VII(a):




VII (a)

where R^1 , R^2 , R^{5a} and R^3 have the same meanings as before, are key intermediates in the preparation of compounds of Formula II.


5 The compound of Formula VII(a) in which R^1 and R^2 represent methyl, R^{5a} represents methoxy, and R^3 represents hydrogen is 5-methoxy-N,N- dimethyltryptamine, available commercially from Sigma, and is the preferred starting material for many of the compounds of Formula II.

10 Other compounds in accordance with Formula VII(a) may be prepared by a variety of routes, such as:



15 where R^{5a} and R^3 have the same meanings as before. The reaction involves the well-known Fischer-indole cyclisation, with in situ substitution of the chlorine by amino (see J. Med. Chem., 1994, 37, 3023). If required, the primary amino group of VIII may be reductively alkylated by reaction with formaldehyde in the presence of sodium cyanoborohydride
 20 to provide the compounds VII(a) in which $R^1 = R^2 = \text{methyl}$.


An alternative route to the compounds VIII is depicted in the following reaction scheme:



5

where R^{5a} and R^3 have the same meanings as before. Step (1) involves coupling of the indole IX with 1-dimethylamino-2-nitroethylene in the presence of trifluoroacetic acid, and step (2) involves reduction with lithium aluminium hydride. Both steps are described in J. Med. Chem.,
 10 1993, 36, 4069.

A further route to the compounds VII(a), also starting from the indoles IX, is depicted in the following reaction scheme:




15

where R^1 , R^2 , R^3 and R^{5a} have the same meanings as before. Step (1) involves reaction of the indoles IX with oxalyl chloride, followed by


condensation with an amine R^1R^2NH . Step (2) involves reduction by lithium aluminium hydride of the α -ketoamide group.

The indoles IX may be prepared by the standard Fischer-indole synthesis, or more conveniently (when R^3 does not represent hydrogen) by
 5 treatment of the N-acyl derivatives X with n-butyllithium:




where R^{5a} is defined as before and R^{3a} represents any group represented by
 10 R^3 except hydrogen (see J. Med. Chem., 1993, 36, 4069).

A preferred route to the compounds of Formula II(b) is shown in the following reaction scheme, in which R^1 , R^2 and Ar have the same meanings as before, and the starting material (5-methoxytryptophol) is commercially available from Sigma.



Step (1) involves reaction of 5-methoxytryptophol with at least two equivalents of arylsulphonyl chloride in the presence of base. Step (2) involves reaction of the resulting arylsulphonate ester with R^1R^2NH so as to cause nucleophilic displacement of the arylsulphonate group.

Representative compounds of Formula III may be prepared by the route shown in the following reaction scheme, in which R^4 has the same meaning as before, and the ketone XI is prepared as described in J. Chem. Soc. Perkin Trans. I, 1972, 1121:



In step (1), reaction of XI with R^4-Cl in the presence of base affords the N-substituted derivative, which is then converted (step (2)) to the α -bromoketone by treatment with phenyl trimethyl ammonium bromide perbromide. Nucleophilic displacement of the bromine atom is effected by reaction with sodium azide in DMF (step (3)), and catalytic hydrogenation of the azide over Pd/C (step (4)) affords the corresponding α -aminoketone. Finally, the ketone is reduced by reaction with sodium borohydride (step

(5)) to provide the aminoalcohol XII as a racemic mixture. If necessary, the primary amino group of XII may be reductively alkylated (e.g. by hydrogenation over Pd/C in the presence of formaldehyde, acetaldehyde etc.) to form the corresponding N,N-dialkyl derivatives. Additionally, or
5 alternatively, the hydroxyl group of XII may be alkylated by the standard methods of ether synthesis.

It will be understood that any compound of Formula I, II or III initially obtained from any of the above processes may, where appropriate, subsequently be elaborated into a further compound of Formula I, II or III
10 respectively by techniques known from the art. For example, a compound of Formula I, II or III in which R^3 represents hydrogen may be reacted with tert-butyllithium to provide the 2-lithio derivative, which may then be reacted with an alkyl, aryl or heteroaryl aldehyde to form a secondary alcohol, which in turn may be oxidised using manganese IV oxide to give
15 the compounds of Formula I, II or III in which R^3 represents alkylcarbonyl, arylcarbonyl or heteroarylcarbonyl. Alternatively, the aforementioned secondary alcohols formed by the reaction of the lithio derivative with an alkyl aldehyde may be subjected to dehydrative elimination to form compounds of Formula I, II or III in which R^3
20 represents alkenyl. Furthermore, compounds of Formula I, II or III in which either or both of R^1 and R^2 represents hydrogen may be alkylated by any of the standard techniques of N-alkylation to provide the corresponding compounds of Formulae I-III in which either or both of R^1 and R^2 represents alkyl. Suitable techniques of N-alkylation include
25 treatment with the appropriate aldehyde or ketone in the presence of sodium cyanoborohydride. Additionally, a compound of Formula I, II or III in which R^5 represents benzyloxy may be converted by catalytic hydrogenation to the corresponding compound of Formula I, II or III in which R^5 represents hydroxy, which in turn may be alkylated by standard
30 methods to provide a corresponding compound in which R^5 represents alkoxy.

Where the above-described processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The novel compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques such as preparative HPLC, or the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-*p*-toluoyl-d-tartaric acid and/or (+)-di-*p*-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The following Examples illustrate the preparation of compounds according to the invention.

EXAMPLE 12-[1-(Benzenesulphonyl)-1*H*-indol-4-yl]ethylamine Hydrogen Oxalate

5

a) Methyl 1-(benzenesulphonyl)-1*H*-indole-4-carboxylate

A mixture of methyl indole-4-carboxylate (*J. Org. Chem.* **1980**, *45*, 3350) (5.0g, 28.5mmol), benzenesulphonyl chloride (10.1g, 57.1mmol) and anhydrous potassium carbonate (15.78g, 114mmol) in butan-3-one (150ml) was refluxed under nitrogen for 15 hours. Additional benzenesulphonyl chloride (10.1g) and anhydrous potassium carbonate (10.1g) were added and refluxing was continued for 5 hours. The mixture was diluted with water (300ml) and extracted with diethyl ether (3 x 300ml). The combined organic solutions were dried (MgSO₄) and concentrated to give a white solid which was recrystallised from ethanol (6.35g, 72%); δ_H (360MHz, CDCl₃), 8.21 (1H, d, J=8.3Hz, Ar-H), 7.96 (1H, dd, J=8.1 and 0.7Hz, Ar-H), 7.87-7.84 (2H, m, Ar-H), 7.69 (1H, d, J=3.7Hz, Ar-H), 7.56-7.33 (5H, m, Ar-H), 3.94 (3H, s, -OMe); *m/z* (CI) 316 (M⁺+1).

1520 b) 1-(Benzenesulphonyl)-1*H*-indole-4-carboxaldehyde

To a cooled (-70°C) and stirred solution of methyl 1-(benzenesulphonyl)-1*H*-indole-4-carboxylate (4.5g, 14.3mmol) in anhydrous diethyl ether (50ml) was added diisobutylaluminium hydride (1M in toluene; 37ml) over 30 minutes, under a nitrogen atmosphere. After being stirred at -70°C for 3 hours, excess diisobutylaluminium hydride was destroyed by dropwise addition of methanol (3ml) and the mixture was allowed to warm to room temperature. 5% Aqueous citric acid (100ml) was added, products were extracted with ethyl acetate (3 x 100ml) and the organic phases were dried (MgSO₄) and concentrated. Flash chromatography of the residue (silica gel, hexane-ethyl acetate,

30

50:50) gave 2.6g (63%) of the intermediate 1-(benzenesulphonyl)-1*H*-indole-4-methanol as a thick colourless oil.

To a solution of the above alcohol (2.5g) in anhydrous dichloromethane (200ml) was added activated manganese (IV) oxide (9g) and the resulting mixture was stirred for 15 hours at room temperature, under a nitrogen atmosphere. The manganese residues were removed by filtration, washed with chloroform and the filtrate was concentrated under vacuum to give 1.96g (79%) of the title compound as a white solid; δ_{H} (250MHz, CDCl_3) 7.42-7.56 (5H, m), 7.72-7.77 (2H, m), 7.86-7.90 (2H, m), 8.28 (1H, d, $J=8.25\text{Hz}$); 10.18 (1H, s); m/z (EI) 285 (M^+). (Found: m/z 285.0444. $\text{C}_{15}\text{H}_{11}\text{NO}_3\text{S}$ requires: m/z 285.0460).

c) 1-(Benzenesulphonyl)-4-(2-nitrovinyl)-1*H*-indole

To a solution of 1-benzenesulphonyl-1*H*-indole-4-carboxaldehyde (1.9g, 6.6mmol) in nitromethane (12ml) was added ammonium acetate (205mg, 2.7mmol) and the resulting mixture was refluxed for 2 hours. Water (50ml) was added and products were extracted with ethyl acetate (2 x 100ml). The combined organic phases were washed with brine (1 x 30ml), dried (MgSO_4) and concentrated. Flash chromatography (silica gel, dichloromethane-hexane, 75:25) of the residue gave 1.95g (90%) of the title compound as a pale yellow solid; δ_{H} (250MHz, CDCl_3) 8.27 (1H, d, $J=13.7\text{Hz}$), 8.14 (1H, d, $J=8.2\text{Hz}$), 7.92-7.87 (2H, m), 7.75 (1H, d, $J=3.7\text{Hz}$), 7.65 (1H, d, $J=13.7\text{Hz}$), 7.62-7.34 (5H, m), 6.89 (1H, dd, $J=3.7$ and 0.9Hz); m/z (CI) 299 (M^++1).

d) 2-[1-(Benzenesulphonyl)-1*H*-indol-4-yl]ethylamine Hydrogen Oxalate

Zinc amalgam was prepared by mixing zinc dust (6g) and mercury (II) chloride (600mg) in a mixture of water (6ml) and concentrated hydrochloric acid (300 μl) and stirring at room temperature for 10 minutes. The aqueous phase was decanted off and the amalgam was then covered with methanol.

To a cooled (0°C) and stirred solution of 1-(benzenesulphonyl)-4-(2-nitrovinyl)-1*H*-indole (1.9g) in a mixture of methanol (35ml) and concentrated hydrochloric acid (20ml) was added portionwise, over 30 minutes, the zinc amalgam prepared above. The mixture was stirred at 5 0°C for a further 30 minutes and at room temperature for 2 hours before the solvent was decanted off and evaporated under vacuum. The remaining residue was partitioned between 1M aqueous ammonia (50ml) and diethyl ether (75ml) and the aqueous phase was re-extracted with diethyl ether (2 x 75ml). The combined ethereal solutions were dried 10 (Na₂SO₄) and concentrated. Flash chromatography of the residue (silica gel, dichloromethane-methanol-ammonia, 40:8:1) gave 950mg of the title compound free base as a thick colourless oil; δ_H (250MHZ, CDCl₃), 7.86 (3H, d), 7.60-7.36 (4H, m), 7.20 (1H, t), 7.04 (1H, d), 6.78 (1H, d), 3.06 (4H, s); m/z (EI) 301 (M⁺+1).

15

EXAMPLE 2

N,N-Dimethyl 2-[1-(benzenesulphonyl)-1*H*-indol-4-yl]ethylamine

Hydrogen oxalate

20 A solution of aqueous formaldehyde (38% w/v; 790 μ l) in methanol (10ml) was added dropwise to a cooled (0°C) and stirred mixture of 2-[1-(benzenesulphonyl)-1*H*-indol-4-yl]ethylamine (750mg), sodium cyanoborohydride (345mg) and glacial acetic acid (715 μ l) in methanol (30ml). The mixture was then allowed to warm to room temperature and 25 it was stirred for 1 hour before saturated aqueous potassium carbonate (20ml) was added and the methanol was removed under vacuum. The remaining residue was diluted with water (20ml) and products were extracted with ethyl acetate (3 x 50ml). The combined organic solutions were washed with brine (1 x 30ml), dried (MgSO₄) and concentrated. 30 Flash chromatography of the residue (silica gel, dichloromethane-methanol, 90:10) gave 550mg (67%) of the title compound as a colourless

thick oil. The oxalate salt was prepared and recrystallised from isopropanol-ether; mp 192-195°C; δ_H (360MHz, D₂O) 7.93-7.89 (3H, m), 7.77 (1H, d, J=3.8Hz), 7.60 (1H, t, J=7.6Hz), 7.50-7.45 (2H, m), 7.35 (1H, t, J=8.2Hz), 7.19 (1H, d, J=7.5Hz), 6.91 (1H, d, J=3.9Hz), 3.43-3.37 (2H, m), 3.28-3.22 (2H, m), 2.89 (6H, s); m/z (EI) 329 (M⁺+1). (Found: C, 56.73; H, 5.26; N, 6.65. C₁₈H₂₀N₂O₂S.1.0 C₂H₂O₄.0.2 H₂O requires: C, 56.91; H, 5.35; N, 6.64%). (Found: m/z 328.1259. C₁₈H₂₀N₂O₂S requires: m/z 328.1246).

10

EXAMPLE 3

N,N-Dimethyl 2-[1-(dimethylamino)sulphonyl-1H-indol-4-yl]ethylamine. Hydrogen Oxalate

The title compound was prepared from 4-[2-(dimethylamino)ethyl]-
1H-indole (P.E. Persons et al., *Eur. J. Med. Chem.*, 1991, 26, 473-475) and
N,N-dimethylaminosulphonyl chloride following a similar method to that
described for Example 6. The oxalate salt was prepared from ethanol-
diethyl ether; mp 153-155°C; δ_H (360MHz, DMSO-d₆) 7.78 (1H, d, J=8.6Hz), 7.70 (1H, d, J=3.6Hz), 7.30 (1H, t, J=7.6Hz), 7.15 (1H, d, J=7.3Hz), 6.98 (1H, d, J=3.6Hz), 3.23 (4H, s), 2.80 (6H, s), 2.79 (6H, s);
 m/z (EI) 296 (M⁺+1). (Found: C, 49.60; H, 5.89; N, 10.35. C₁₄H₂₁N₃O₂S.1.1 C₂H₂O₄ requires: C, 49.33; H, 5.93; N, 10.65%).

25

EXAMPLE 4

N,N-Dimethyl 3-[1-(benzenesulphonyl)-1H-indol-4-yl]propylamine. Hydrogen Oxalate

a) 3-[1-(Benzenesulphonyl)-1H-indol-4-yl]acrylic acid ethyl ester

A solution of 1-(benzenesulphonyl)-1H-indole-4-carboxaldehyde (2.0 g, 7 mmol) and (carbethoxymethylene)triphenylphosphorane (2.69 g, 7.7

mmol) in anhydrous dichloromethane (50 ml) was heated to reflux for 3 h under a nitrogen atmosphere. The reaction was concentrated and purified by dry flash chromatography (dichloromethane) to give the title compound as a colourless solid (2.49 g, 100%). δ_{H} (360MHz, CDCl_3), 1.34 (3H, t, $J=7.2\text{Hz}$), 4.27 (2H, q, $J=7.2\text{Hz}$), 6.51 (1H, d, $J=17.2\text{Hz}$), 6.94 (1H, d, $J=0.72\text{Hz}$), 7.32 (1H, t, $J=7.9\text{Hz}$), 7.43-7.67 (4H, m), 7.67 (1H, d, $J=3.6\text{Hz}$), 7.68-7.89 (2H, m), 7.94-8.03 (2H, m).

b) 3-[1-(Benzenesulphonyl)-1*H*-indol-4-yl]propionic acid ethyl ester

3-[1-(Benzenesulphonyl)-1*H*-indol-4-yl]acrylic acid ethyl ester (2.48 g, 6.98 mmol) was hydrogenated over 10% Pd-C (500 mg) in ethyl acetate-ethanol (4:1; 75 ml), under an atmosphere of hydrogen. The catalyst was removed by filtration after 4 h, and the solvents were removed to obtain the title compound as a colourless oil which crystallised on standing (2.47 g, 100%); δ_{H} (CDCl_3 , 360MHz) 1.10 (3H, t, $J=6.8\text{Hz}$), 2.57 (2H, t, $J=7.5\text{Hz}$), 3.06 (2H, t, $J=7.5\text{Hz}$), 4.02 (2H, q, $J=7.2\text{Hz}$), 6.66 (1H, d, $J=0.72\text{Hz}$), 6.98 (1H, d, $J=7.5\text{Hz}$), 7.14-7.18 (1H, m), 7.34-7.38 (2H, m), 7.43-7.48 (1H, m), 7.50 (1H, d, $J=3.96\text{Hz}$), 7.77-7.82 (3H, m).

c) *N,N*-Dimethyl-3-[1-(benzenesulphonyl)-1*H*-indol-4-yl]propionamide

Trimethylaluminium (1.4 ml, 2M solution in hexanes) was added to a suspension of dimethylamine hydrochloride (228 mg, 2.8 mmol) in 10 ml benzene. The reaction was allowed to stir for 1h before dropwise addition of a solution of 3-[1-(benzenesulphonyl)-1*H*-indol-4-yl]propionic acid ethyl ester (0.5g, 1.4 mmol) in benzene (10 ml). The reaction was heated at reflux for 18 h, allowed to cool and then quenched by addition of HCl (2N, 5 ml). The reaction was partitioned between H_2O -EtOAc. The organic phase was washed with brine, dried (MgSO_4), and concentrated to give the title compound as an oil which crystallised (527 mg, 100%); δ_{H} (CDCl_3 , 250MHz), 2.63 (2H, t, $J=7.5\text{Hz}$), 2.79 (2H, br s), 3.16 (2H, t, $J=8.5\text{Hz}$), 6.74

(1H, d, $J=0.7\text{Hz}$), 6.08 (1H, d, $J=7.5\text{Hz}$), 7.21-7.27 (1H, m), 7.41-7.58 (4H, m), 7.84-7.90 (3H, m).

d) *N,N*-Dimethyl 3-[1-(benzenesulphonyl)-1*H*-indol-4-yl]propylamine

5 Hydrogen Oxalate

A solution of the product from above (527 mg, 1.4 mmol) in 15 ml THF was treated dropwise with lithium aluminium hydride (1.4 ml, 1.0M THF) whilst stirring under a nitrogen atmosphere. The reaction was stirred for 5 h, and quenched with water (1 ml). The reaction was filtered
10 through a glass fibre pad, and the filtrate concentrated. The residue was partitioned between 2N NaOH-EtOAc. The organic phase was dried (Na_2SO_4), concentrated and purified by chromatography [CH_2Cl_2 -MeOH- NH_4OH (95:5:0.5)] to give the title compound free base as an oil (2.75 mg, 50%). The oxalate salt was prepared in diethyl ether; mp 142-144°C.
15 (Found: C, 56.29; H, 5.06; N, 6.12. $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}_3\text{S} \cdot 1.3 \text{ C}_2\text{H}_2\text{O}_4$ requires C, 56.46; H, 5.40; N, 6.10). δ_{H} (d_6 -DMSO, 250MHz) 1.8-2.0 (2H, m), 2.67 (6H, s), 2.78-2.84 (2H, m), 2.98-3.04 (2H, m), 6.97 (1H, d, $J=3.75\text{Hz}$), 7.11 (1H, d, $J=7\text{Hz}$), 7.29 (1H, t, $J=7.5\text{Hz}$), 7.11 (1H, d, $J=7\text{Hz}$), 7.29 (1H, t, $J=7.5\text{Hz}$), 7.57-7.73 (3H, m), 7.79-7.85 (2H, m), 7.98-9.01 (2H, m); m/z
20 (ES⁺) 343 (M^++1).

EXAMPLE 5

N,N-Dimethyl 2-[1-(benzenesulphonyl)-2-benzoyl-1*H*-indol-4-yl]ethylamine. Hydrogen oxalate
25

To a cooled (-78°C) and stirred solution of *N,N*-dimethyl 2-[1-(benzenesulphonyl)-1*H*-indol-4-yl]ethylamine (1.16 g, 3.6 mmol) in anhydrous tetrahydrofuran (8 ml) was added dropwise, under nitrogen, *tert*-butyllithium (1.7 M in pentane, 25 ml). The resulting solution was
30 stirred at -78°C for 30 minutes and then it was allowed to warm to 0°C over 1 h. After being re-cooled to -78°C, a solution of freshly distilled

benzaldehyde (415 mg, 3.9 mmol) in anhydrous tetrahydrofuran (4 ml) was added and the mixture was allowed to warm to room temperature over 3 h. The reaction was quenched by addition of water (20 ml) and products were extracted with ethyl acetate (2 x 75ml), dried (MgSO₄) and concentrated. Flash chromatography (silica gel, dichloromethane-ethanol, 90:10 to dichloromethane-methanol-ammonia, 60:8:1) of the residue gave 600mg of [1-(benzenesulphonyl)-4-[2-(dimethylamino)ethyl]-1*H*-indol-2-yl]phenylmethanol as a white solid.

To a solution of the above alcohol (350 mg) in anhydrous dichloromethane (20 ml) was added activated manganese (IV) oxide (250 mg) and the resulting mixture was stirred vigorously at room temperature for 30 minutes. The manganese residues were filtered off, washed with dichloromethane and the filtrate was concentrated under vacuum. Flash chromatography of the residual oil (silica gel, dichloromethane-ethanol, 90:10 to dichloromethane-methanol-ammonia, 60:8:1) gave 330 mg of the title compound as a yellow oil. The hydrogen oxalate salt was prepared and recrystallised from isopropanol-diethyl ether; mp 154-156°C; δ_H (360MHz, d₆-DMSO) 8.03-7.93 (5H, m), 7.78-7.72 (2H, m), 7.68-7.59 (4H, m), 7.51-7.46 (2H, m), 7.26 (1H, d, *J*=7.4Hz), 3.16 (4H, s, -CH₂CH₂-), 2.73 (6H, s, -NMe₂). (Found: C, 61.70; H, 5.07; N, 5.33. C₂₅H₂₄N₂O₃S.1.0 C₂H₂O₄.0.2 H₂O requires: C, 61.66; H, 5.02; N, 5.33%).

EXAMPLE 6

N,N-Dimethyl 2-[1-(benzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine Hydrochloride

To a stirred two-phase mixture of 5-methoxy-*N,N*-dimethyl-tryptamine (2 g, 9.16 mmol) and tetrabutylammonium hydrogen sulphate (311 mg, 0.916 mmol) in dichloromethane (120 ml) and 50% aqueous sodium hydroxide (30 ml) was added benzenesulphonyl chloride (2.43 g, 13.74 mmol) over 30 minutes. The mixture was stirred at room

temperature for 20 minutes, diluted with ethyl acetate (250 ml) and water (125 ml), and the organic phase was decanted off, washed with brine (2 x 65 ml), dried (MgSO₄) and concentrated. Flash chromatography (silica gel, dichloromethane-methanol-ammonia, 95:5:0.5) of the residue afforded
5 3.166 g (96%) of the title compound free base as a thick oil which solidified on standing. The hydrogen chloride salt was prepared; mp 200-215°C (EtOH); δ_H (360MHz, DMSO-d₆) 10.82 (1H, br s), 7.92 (2H, dd, J=8.6 and 1.3Hz), 7.82 (1H, d, J=9.0Hz), 7.72-7.65 (2H, m), 7.58 (2H, t, J=7.5Hz), 7.26 (1H, d, J=2.5Hz), 6.98 (1H, dd, J=9.0 and 2.5Hz), 3.80 (3H, s), 3.36-
10 3.28 (2H, m), 3.16-3.28 (2H, m), 3.16-3.06 (2H, m), 2.81 (6H, s); m/z (ES) 359 (M⁺⁺¹). (Found: C, 58.04; H, 5.97; N, 7.37. C₁₉H₂₂N₂SO₃.HCl requires: C, 57.79; H, 5.87; N, 7.09%).

EXAMPLE 7

15

N,N-Dimethyl 2-[5-methoxy-1-(4-methylbenzenesulphonyl)-1H-indol-3-yl]ethylamine Hydrochloride

Following a similar procedure to that described in Example 6, but using *p*-toluenesulphonyl chloride instead of benzenesulphonyl chloride,
20 the title compound was prepared in 72% yield as a white solid; mp 214-215°C; δ_H (360 MHz, *d*₆-DMSO) 2.32 (3H, s), 2.81 (6H, s), 3.08 (2H, m), 3.37 (2H, m), 3.79 (3H, s), 6.97 (1H, dd, J=9.0, 2.5 Hz), 7.23 (1H, d, J=2.4 Hz), 7.37 (2H, d, J=8.1 Hz), 7.67 (1H, s), 7.79 (2H, d, J=8.4 Hz), 7.80 (1H, d, J=9.1 Hz), 10.60 (1H, br s); m/z (ES⁺) 373 (MH⁺). Found: C, 57.66; H,
25 6.12; N, 6.74. C₂₀H₂₄N₂O₃S.HCl.0.4H₂O requires C, 57.72; H, 6.25; N, 6.73%.

EXAMPLE 8*N,N*-Dimethyl 2-[1-(4-chlorobenzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine Hydrochloride

5 Following a similar procedure to that described in Example 6, but using 4-chlorobenzenesulphonyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 39% yield as a white solid; mp 222-224°C; δ_{H} (360 MHz, d_6 -DMSO) 2.82 (6H, s), 3.09 (2H, m), 3.35 (2H, m), 3.81 (3H, s), 6.99 (1H, dd, $J=9.0, 2.5$ Hz), 7.27 (1H, d, $J=2.4$ Hz),
10 7.67 (2H, d, $J=8.8$ Hz), 7.70 (1H, s), 7.82 (1H, d, $J=9.0$ Hz), 7.94 (2H, d, $J=8.7$ Hz), 10.65 (1H, br s); m/z (ES^+) 395/393 (MH^+). Found: C, 52.76; H, 4.93; N, 6.39. $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_3\text{S} \cdot \text{HCl} \cdot 0.2\text{H}_2\text{O}$ requires C, 52.71; H, 5.22; N, 6.47%.

15

EXAMPLE 9*N,N*-Dimethyl 2-[1-(3-chlorobenzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine Hydrochloride

20 Following a similar procedure to that described in Example 6, but using 3-chlorobenzenesulphonyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 67% yield as a white solid; mp 191-194 °C; δ_{H} (360 MHz, d_6 -DMSO) 2.81 (6H, s), 3.09 (2H, m), 3.34 (2H, m), 3.80 (3H, s), 7.00 (1H, dd, $J=9.0, 2.5$ Hz), 7.26 (1H, d, $J=2.4$ Hz),
25 7.61 (1H, t, $J=8.0$ Hz), 7.73 (1H, s), 7.77 (1H, dt), 7.84 (1H, d, $J=9.1$ Hz), 7.89 (1H, dt), 7.97 (1H, t, $J=1.9$ Hz), 10.59 (1H, br s); m/z (ES^+) 395/393 (MH^+). Found: C, 52.88; H, 5.12; N, 6.62. $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_3\text{S} \cdot \text{HCl}$ requires C, 53.15; H, 5.16; N, 6.52%.

EXAMPLE 10*N,N*-Dimethyl 2-[5-methoxy-1-(2-naphthalenesulphonyl)-1*H*-indol-3-yl]ethylamine Hydrochloride

5 Following a similar procedure to that described in Example 6, but using 2-naphthalenesulphonyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 56% yield as a white solid; mp 191-193 °C; δ_{H} (360 MHz, d_6 -DMSO) 2.81 (6H, s), 3.09 (2H, m), 3.35 (2H, m), 3.78 (3H, s), 6.98 (1H, dd, $J=9.0, 2.4$ Hz), 7.23 (1H, d, $J=2.4$ Hz),
10 7.72 (2H, m), 7.78 (1H, s), 7.81 (1H, dd, $J=8.8, 1.9$ Hz), 7.90 (1H, d, $J=9.1$ Hz), 8.01 (1H, d, $J=7.7$ Hz), 8.08 (1H, d, $J=8.8$ Hz), 8.21 (1H, d), 8.76 (1H, s), 10.60 (1H, br s); m/z (ES⁺) 409 (MH⁺). Found: C, 62.19; H, 5.51; N, 6.41. C₂₃H₂₄N₂O₃S.HCl requires C, 62.08; H, 5.66; N, 6.30%.

15

EXAMPLE 11*N,N*-Dimethyl 2-[5-methoxy-1-(4-methoxybenzenesulphonyl)-1*H*-indol-3-yl]ethylamine Hydrochloride

20 Following a similar procedure to that described in Example 6, but using 4-methoxybenzenesulphonyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 61% yield as a white solid; mp 152-155°C; δ_{H} (360 MHz, d_6 -DMSO) 2.82 (6H, s), 3.08 (2H, m), 3.34 (2H, m), 3.80 (3H, s), 3.81 (3H, s), 6.98 (1H, dd, $J=9.0, 2.5$ Hz), 7.08 (2H, d, $J=9.0$ Hz), 7.24 (1H, d, $J=2.4$ Hz), 7.67 (1H, s), 7.81 (1H, d, $J=9.0$ Hz), 7.86
25 (2H, d, $J=9.1$ Hz), 10.55 (1H, br s); m/z (ES⁺) 389 (MH⁺). Found: C, 56.17; H, 5.94; N, 6.63. C₂₀H₂₄N₂O₄S.HCl requires C, 56.53; H, 5.93; N, 6.59%.

EXAMPLE 12*N,N*-Dimethyl 2-[1-(2-chlorobenzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine Hydrochloride

5 Following a similar procedure to that described in Example 6, but using 2-chlorobenzenesulphonyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 43% yield as a white solid; mp 185-188°C; δ_{H} (360 MHz, d_6 -DMSO) 2.84 (6H, s), 3.14 (2H, m), 3.37 (2H, m), 3.81 (3H, s), 6.93 (1H, dd, $J=9.0, 2.4$ Hz), 7.31 (1H, d, $J=2.4$ Hz),
10 7.54 (1H, d, $J=9.0$ Hz), 7.63 (1H, td), 7.68 (1H, dd), 7.74 (1H, td), 7.78 (1H, s), 8.14 (1H, dd, $J=7.9, 1.5$ Hz), 10.45 (1H, br s); m/z (ES⁺) 395/393 (MH⁺). Found: C, 53.09; H, 5.06; N, 6.47. C₁₉H₂₁ClN₂O₃S.HCl requires C, 53.15; H, 5.16; N, 6.52%.

15

EXAMPLE 13*N,N*-Dimethyl 2-(1-benzoyl-5-methoxy-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

20 Following a similar procedure to that described in Example 6, but using benzoyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 53% yield as a white solid; mp 199-201°C; δ_{H} (360 MHz, d_6 -DMSO) 2.78 (6H, s), 3.05 (2H, m), 3.27 (2H, m), 3.86 (3H, s), 7.02 (1H, dd, $J=9.0, 2.4$ Hz), 7.28 (1H, d, $J=2.4$ Hz), 7.32 (1H, s), 7.61 (2H, t, $J=7.0$ Hz), 7.70 (1H, t, $J=7.3$ Hz), 7.75 (2H, d, $J=6.9$ Hz), 8.19 (1H, d,
25 $J=9.0$ Hz); m/z (ES⁺) 323 (MH⁺). Found: C, 63.89; H, 5.80; N, 7.10. C₂₀H₂₂N₂O₂.C₂H₂O₄ requires C, 64.07; H, 5.87; N, 6.79%.

EXAMPLE 14*N,N*-Dimethyl 2-[5-methoxy-1-(2-thiophenesulphonyl)-1*H*-indol-3-yl]ethylamine Hydrogen Oxalate

5 Following a similar procedure to that described in Example 6, but using 2-thiophenesulphonyl chloride instead of benzenesulphonyl chloride, the title compound was prepared in 25% yield as a white solid; mp 215-220°C; δ_{H} (360 MHz, *d*₆-DMSO) 2.78 (6H, s), 3.03 (2H, m), 3.29 (2H, m), 3.81 (3H, s), 7.16 (1H, m), 7.20 (1H, d, *J*=2.3 Hz), 7.61 (1H, s), 7.81 (1H, d, *J*=9.2 Hz), 7.83 (1H, m), 8.01 (1H, dd, *J*=5.1, 1.3 Hz); *m/z* (ES⁺) 365 (MH⁺).
10 Found: C, 50.45; H, 4.89; N, 6.27. C₁₇H₂₀N₂O₃S₂.C₂H₂O₄ requires C, 50.21; H, 4.88; N, 6.16%.

EXAMPLE 15

15

2-[1-(Benzenesulphonyl)-5-methoxy-1*H*-indol-3-yl]ethylamine Hydrochloride

 To a stirred solution of 5-methoxytryptamine (3.5 g, 18.4 mmol) in dichloromethane (100 ml) was added dropwise a solution of di-*tert*-butyl dicarbonate (4.02 g, 18.4 mmol) in dichloromethane (25 ml) over 5
20 minutes. The resulting pale brown solution was allowed to stand at room temperature for 25 h before solvents were removed under vacuum to give a thick oil.

 To a stirred two-phase mixture of a solution of the above 3-[2-(*tert*-
25 butyloxycarbonylamino)ethyl]-5-methoxy-1*H*-indole in dichloromethane (200 ml) and 50% aqueous sodium hydroxide (50 ml) was added tetrabutylammonium hydrogen sulphate (611 mg, 1.8 mmol) and the mixture was stirred for 10 minutes. Benzenesulphonyl chloride (3.52 ml, 27.6 mmol) was added dropwise *via* syringe and the mixture was
30 vigorously stirred for 30 minutes. The mixture was diluted with diethyl ether (500 ml), washed with water (100 ml), brine (2 x 70 ml), dried

(MgSO₄) and concentrated. Flash chromatography of the residue (silica gel, dichloromethane-diethyl ether, 90:10) followed by trituration with hexane-diethyl ether (10:1; 100 ml) afforded 1-(benzenesulphonyl)-3-[2-(*tert*-butyloxycarbonylamino)ethyl]-5-methoxy-1*H*-indole (7.55 g, 95%) as a white solid; δ_H (360MHz, CDCl₃) 7.88 (1H, d, J=8.8Hz), 7.84-7.80 (2H, m), 7.54-7.48 (1H, m), 7.45-7.38 (2H, m), 7.32 (1H, s), 6.96-6.90 (2H, m), 4.55 (1H, br s), 3.82 (3H, s), 3.44-3.36 (2H, m), 2.82 (2H, t, J=6.8Hz), 1.45 (9H, s).

A solution of the above protected tryptamine (2.1 g, in a mixture of dichloromethane and trifluoroacetic acid (1:1; 60 ml) was allowed to stand at room temperature for 3 h. The solvents were removed under vacuum and the residue was azeotroped with toluene-methanol (5:1; 100 ml). The residue was dissolved in 4N sodium hydroxide solution (50 ml) and extracted with ethyl acetate (3 x 100 ml). The combined organic solutions were washed with brine (1 x 50 ml), dried (Na₂SO₄) and concentrated. The hydrochloride salt was prepared and recrystallised from acetonitrile to give 1.53 g of the title compound; mp 117-120°C; δ_H (360MHz, D₂O) 7.81 (2H, d, J=7.5Hz), 7.74 (1H, d, J=9.0Hz), 7.56 (1H, s), 7.54 (1H, t, J=7.5Hz), 7.41 (2H, t, J=8.1Hz), 7.00 (1H, d, J=2.3Hz), 6.89 (1H, dd, J=9.0 and 2.3Hz), 6.89 (1H, dd, J=9.0 and 2.3Hz), 3.78 (3H, s), 3.29 (2H, t, J=7.6Hz), 3.01 (2H, t, J=7.6Hz); *m/z* (ES) 331 (M⁺+1). (Found: C, 54.07; H, 5.39; N, 7.41. C₁₇H₁₈N₂O₃S.1.0 HCl.0.6 H₂O requires: C, 54.06; H, 5.39; N, 7.42%).

EXAMPLE 16

1-Benzenesulphonyl-5-methoxy-3-[(2-pyrrolidin-1-yl)ethyl]-1*H*-indole
Hydrogen Oxalate

a) Benzenesulphonic acid 2-(1-benzenesulphonyl-5-methoxy-1*H*-indol-3-yl)ethyl ester

To a stirred two-phase mixture of 5-methoxytryptophol (0.150 g, 0.785 mmol) and tetrabutylammonium hydrogen sulphate (27 mg, 0.0785 mmol) in 50% aqueous NaOH (3 mL) and dichloromethane (30 mL) was added benzenesulphonyl chloride (0.25 mL, 1.96 mmol) dropwise over 5 min. The mixture was stirred for 20 min, more benzenesulphonyl chloride 0.10 mL, 0.784 mmol) was added dropwise, and the mixture was stirred for a further hour. The organic layer was washed with brine, dried (Na₂SO₄) and evaporated in vacuo. The residue was purified by flash chromatography (silica gel, 50-100% EtOAc/hexane) to afford 0.095 g (37%) of 2-(1-benzenesulphonyl-5-methoxy-1*H*-indol-3-yl)ethanol and 0.191 g (52%) of the title compound; δ_{H} (360 MHz, *d*₆-DMSO) 2.98 (2H, t, *J* 6.1 Hz), 3.71 (3H, s), 4.30 (2H, t, *J*=6.2 Hz), 6.92 (1H, dd, *J*=8.9, 2.5 Hz), 6.95 (1H, d, *J*=2.2 Hz), 7.47 (2H, t, *J*=7.8 Hz), 7.53-7.59 (3H, m), 7.63-7.67 (4H, m), 7.77 (1H, d, *J*=8.9 Hz), 7.89 (2H, m); *m/z* (ES⁺) 510 (M+K)⁺, 494 (M+Na)⁺, 489 (M+NH₄)⁺, 472 (M+H)⁺.

b) 1-Benzenesulphonyl-5-methoxy-3-[(2-pyrrolidin-1-yl)ethyl]-1*H*-indole
Hydrogen Oxalate

To a stirred solution of benzenesulphonic acid 2-(1-benzenesulphonyl-5-methoxy-1*H*-indol-3-yl)ethyl ester (0.191 g, 0.405 mmol) in anhydrous 2-propanol (10 mL) under nitrogen was added anhydrous potassium carbonate (0.112 g, 0.810 mmol), then a solution of pyrrolidine (0.17 mL, 2.03 mmol) in anhydrous 2-propanol (3 mL). The mixture was heated at reflux for 4 h, and the solvents were then removed in vacuo. The residue was partitioned between dichloromethane and water, and the aqueous layer was extracted further with dichloromethane. The combined organic extracts were washed with brine (2 x 25 mL), dried (Na₂SO₄), and evaporated in vacuo. The residue was purified by flash chromatography (silica gel, CH₂Cl₂-MeOH-NH₃; 95:5:0.5) to give 0.102 g

(65%) of the title compound free base. The oxalate salt was prepared in EtOH-Et₂O as a white solid; mp 217-220°C; δ_{H} (360 MHz, *d*₆-DMSO) 1.94 (4H, s), 3.03 (2H, m), 3.27 (4H, s), 3.40 (2H, m), 3.79 (3H, s), 6.98 (1H, dd, *J*=9.0, 2.3 Hz), 7.17 (1H, d, *J*=2.3 Hz), 7.58 (2H, t, *J*=8.0 Hz), 7.69 (2H, m), 7.82 (1H, d, *J*=9.0 Hz), 7.92 (2H, d, *J*=7.9 Hz); *m/z* (ES⁺) 385 (MH⁺).
Found: C, 57.83; H, 5.31; N, 5.87. C₂₁H₂₄N₂O₃S.C₂H₂O₄ requires C, 58.22; H, 5.52; N, 5.90%.

EXAMPLE 17

10

1-Benzenesulphonyl-5-methoxy-3-[(2-piperidin-1-yl)ethyl]-1*H*-indole Hydrogen Oxalate

Following a similar procedure to that described in Example 16, step b, but using piperidine instead of pyrrolidine, the title compound was prepared in 90% yield as a white solid; mp 231-234°C; δ_{H} (360 MHz, *d*₆-DMSO) 1.55 (2H, m), 1.75 (4H, m), 3.04 (2H, m), 3.15 (4H, s), 3.27 (2H, m), 3.79 (3H, s), 6.98 (1H, dd, *J*=9.0, 2.4 Hz), 7.15 (1H, d, *J*=2.4 Hz), 7.58 (2H, t, *J*=8.0 Hz), 7.68 (2H, m), 7.82 (1H, d, *J*=9.0 Hz), 7.91 (2H, d, *J*=7.7 Hz); *m/z* (ES⁺) 399 (MH⁺). Found: C, 58.40; H, 5.91; N, 5.81.
C₂₂H₂₆N₂O₃S.C₂H₂O₄.0.3H₂O requires C, 58.36; H, 5.84; N, 5.67%.

20

EXAMPLE 18

1-Benzenesulphonyl-5-methoxy-3-[(2-piperazin-1-yl)ethyl]-1*H*-indole Hydrogen Oxalate

25

Following a similar procedure to that described in Example 16, step b, but using piperazine instead of pyrrolidine, the title compound was prepared in 76% yield as a white solid mp 230-232°C; δ_{H} (360 MHz, *d*₆-DMSO) 2.70 (3H, s), 2.82 (8H, br s), 3.09 (4H, br s), 3.78 (3H, s), 6.96 (1H, dd, *J*=9.0 Hz), 7.08 (1H, d), 7.57 (2H, t, *J*=7.5 Hz), 7.60 (1H, s), 7.68 (1H, t, *J*=7.5 Hz), 7.81 (1H, d, *J*=9.0 Hz), 7.90 (2H, d, *J*=7.6 Hz); *m/z* (ES⁺) 414

30

(MH⁺). Found: C, 44.48; H, 4.85; N, 5.13. C₂₂H₂₇N₃O₃S.4C₂H₂O₄.2H₂O requires C, 44.50; H, 4.86; N, 5.19%.

EXAMPLE 19

5

N,N-Dimethyl 2-(1-benzenesulphonyl-5-methoxy-2-methyl-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

a) *N,N*-Dimethyl 2-(5-methoxy-2-methyl-1*H*-indol-3-yl)ethylamine

10 To a stirred solution of 2-(5-methoxy-2-methyl-1*H*-indol-3-yl)ethylamine hydrochloride (Spadoni, G. et al. *J. Med. Chem.* **1993**, *36*, 4069-4074) (1.6 g, 6.55 mmol) in methanol (100 ml) was added a 30% w/v solution of sodium methoxide in methanol (1.24 ml, 6.55 mmol), then acetic acid (1.50 ml, 26.2 mmol), followed by sodium cyanoborohydride
15 (0.82 g, 13.1 mmol). A 38% w/v aqueous solution of formaldehyde (1.24 ml, 15.7 mmol) in methanol (30 ml) was then added dropwise over 40 min, and the mixture was stirred overnight before 4N aqueous NaOH (50 ml) was added. The methanol was removed in vacuo and the aqueous residue was extracted with ethyl acetate. The organic extract was washed with brine
20 (50 ml), dried (Na₂SO₄), and evaporated in vacuo to give 1.215 g (80%) of the title compound; δ_H (360 MHz, CDCl₃) 2.35 (3H, s), 2.36 (6H, s), 2.50 (2H, m), 3.85 (2H, m), 3.85 (3H, s), 6.76 (1H, dd, *J*=8.7, 2.5 Hz), 6.97 (1H, d, *J*=2.3 Hz), 7.13 (1H, d, *J*=8.5 Hz), 7.72 (1H, br s).

25 b) *N,N*-Dimethyl 2-(1-benzenesulphonyl-5-methoxy-2-methyl-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

Following a similar procedure to that described in Example 6, but using *N,N*-dimethyl 2-(5-methoxy-2-methyl-1*H*-indol-3-yl)ethylamine instead of 5-methoxy-*N,N*-dimethyltryptamine the title compound was
30 prepared as a white solid; mp 194-196°C; δ_H (360 MHz, *d*₆-DMSO) 2.54 (3H, s), 2.76 (6H, s), 2.98 (2H, m), 3.44 (2H, m), 3.80 (3H, s), 6.92 (1H, dd,

$J=9.0, 2.5$ Hz), 7.09 (1H, d, $J=2.5$ Hz), 7.56 (2H, t, $J=7.5$ Hz), 7.67 (1H, t, $J=7.5$ Hz), 7.77 (2H, d, $J=7.4$ Hz), 7.94 (1H, d, $J=9.1$ Hz); m/z (ES⁺) 373 (MH⁺). Found: C, 56.16; H, 5.52; N, 6.09. C₂₀H₂₄N₂O₃S.C₂H₂O₄.0.4H₂O requires C, 56.25; H, 5.75; N, 5.96%.

5

Examples 20, 21 and 22 were prepared from the appropriate tryptamine compounds and the corresponding sulphonyl chloride following similar methods to that described for Example 6.

10

EXAMPLE 20

N,N-Dimethyl 2-(1-Benzenesulphonyl-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

mp 196-198°C (EtOH-Et₂O); δ_H (360MHz, DMSO-d₆) 7.97-7.91 (3H, m), 7.74 (1H, s), 7.72-7.64 (2H, m), 7.61 (2H, t, $J=8.0$ Hz), 7.38 (1H, t, $J=7.3$ Hz), 7.29 (1H, t, $J=7.3$ Hz), 3.34-3.26 (2H, m), 7.08-3.00 (2H, m), 2.77 (6H, s); m/z (CI) 329 (M⁺+1). (Found: C, 57.06; 5.40, N, 6.58. C₁₈H₂₀N₂O₂S.1.0 C₂H₂O₄ requires: C, 57.40; H, 5.30; N, 6.69%).

20

EXAMPLE 21

N,N-Dimethyl 2-(1-methylsulphonyl-1*H*-indol-3-yl)ethylamine. Hydrogen Oxalate

mp 191-193°C (EtOH-Et₂O); δ_H (360MHz, DMSO-d₆) 7.83 (1H, d, $J=8.1$ Hz), 7.75 (1H, d, $J=7.3$ Hz), 7.52 (1H, s), 7.44-7.30 (2H, m), 3.37 (3H, s), 3.36-3.28 (2H, m), 3.14-3.05 (2H, m), 2.79 (6H, s); m/z (CI) 267 (M⁺+1). (Found: C, 49.93; H, 5.52; N, 7.76. C₁₃H₁₈N₂O₂S.1.1 C₂H₂O₄ requires: C, 49.96; H, 5.57; N, 7.67%).

EXAMPLE 22*N,N*-Dimethyl 2-(5-methoxy-1-methylsulphonyl-1*H*-indol-3-yl)ethylamine.
Hydrogen Oxalate

5 mp 201-203°C (EtOH-Et₂O); δ_H (360MHz, DMSO-d₆) 7.71 (1H, d, J=9.0Hz), 7.48 (1H, s), 7.25 (1H, d, J=2.4Hz), 7.01 (1H, dd, J=9.0 and 2.4Hz), 3.83 (3H, s), 3.36-3.28 (5H, m and s), 3.10-3.02 (2H, m), 2.79 (6H, s); *m/z* (CI) 297 (M⁺+1). (Found: C, 49.74; H, 5.56; N, 7.07. C₁₄H₂₀N₂O₃S.1.0 C₂H₂O₄ requires: C, 49.73; H, 5.74; N, 7.25%).

10

EXAMPLE 23[3-(2-Dimethylamino-ethyl)-5-hydroxy-1*H*-indol-1-yl]phenylmethanone15 a) 2-(5-Benzoyloxy-1*H*-indol-3-yl)-*N,N*-dimethyl-2-oxo-acetamide

Oxalyl chloride (3.9 ml, 45 mmol) was added to a stirred solution of 5-benzoyloxyindole (5.0g, 22 mmol) in dry tetrahydrofuran (200 mL), at 0°C under nitrogen. The cooling bath was removed and the reaction was stirred at room temperature for 3 hours. The reaction was then cooled to 20 0°C and dimethylamine was bubbled through the solution for 5 minutes (a heavy white precipitate forms). The whole mixture was then poured into 4N hydrochloric acid (50 ml). The layers were separated and the aqueous was extracted with diethylether (x1). The combined organic layers were dried (Na₂SO₄), filtered and evaporated to give the title compound (6.4g, 25 89%) as an off-white solid; δ_H (360 MHz, *d*₆-DMSO) 2.90 (3H, s), 2.98 (3H, s), 5.14 (2H, s), 6.99 (1H, dd, J = 8.3 and 2.0 Hz), 7.31-7.50 (6H, m), 7.71 (1H, d, J = 2.0 Hz), 8.02 (1H, s), 12.2 (1H, br s).

b) [2-(5-Benzoyloxy-1*H*-indol-3-yl)ethyl]dimethylamine

30 A solution of lithium aluminium hydride (1.0M, 120 ml, 120 mmol) was added to a stirred solution of 2-(5-benzoyloxy-1*H*-indol-3-yl)-*N,N*-

dimethyl-2-oxo-acetamide (6.4g, 19.9 mmol) in dry tetrahydrofuran (100 ml) at room temperature under nitrogen. (Gas evolution and precipitate formation were observed) The reaction was then stirred and heated at reflux for 4 hours. The mixture was then cooled to 0°C, and quenched with water (4.5 ml), 4N sodium hydroxide (4.5 ml). Ethyl acetate and anhydrous sodium sulphate were then added. After stirring at room temperature overnight, the precipitate was removed by filtration (washing with ethyl acetate). The filtrate was evaporated and the residue was purified by chromatography on silica gel, eluting with dichloromethane/methanol/aqueous ammonia (80:8:1) to give the title amine (4.0g, 68%) as a thick oil which solidifies upon standing; δ_H (360 MHz, $CDCl_3$) 2.33 (6H, s), 2.58-2.62 (2H, m), 2.87-2.91 (2H, m), 5.10 (2H, s), 6.92 (1H, dd, $J=8.8$ and 2.4 Hz), 6.98 (1H, d, $J=2.2$ Hz), 7.12 (1H, d, $J=2.4$ Hz), 7.23-7.48 (6H, m), 8.0 (1H, br s).

c) [5-Benzoyloxy-3-(2-dimethylamino-ethyl)-1*H*-indol-1-yl]phenylmethanone

Benzoyl chloride (217 ml, 1.9 mmol) was added to a stirred solution of [2-(5-benzyloxy-1*H*-indol-3-yl)ethyl]dimethylamine (500 mg, 1.7 mmol), triethylamine (473 mL, 3.4 mmol) and 4-(dimethylamino)pyridine (20 mg, 0.2 mmol) in dry dichloromethane (5 ml) at 0°C. The reaction was allowed to warm slowly to ambient temperature, and stirred at this temperature overnight. The reaction was quenched with saturated aqueous sodium hydrogen carbonate. The aqueous was then extracted with dichloromethane (x3). The combined extracts were dried (Na_2SO_4), filtered and evaporated. The residue was purified by chromatography on silica gel, eluting with dichloromethane/methanol/aqueous ammonia (80:8:1) to give the title compound (327 mg, 48%) as a thick oil. δ_H (250 MHz, $CDCl_3$) 2.30 (6H, s), 2.52-2.60 (2H, m), 2.78-2.84 (2H, m), 5.16 (2H, s), 7.06-7.10 (3H, m), 7.30-7.64 (10H, m), 8.28 (0.5H, s), 8.32 (0.5H, s).

d) [3-(2-Dimethylamino-ethyl)-5-hydroxy-1*H*-indol-1-yl]phenylmethanone

A solution of [5-benzyloxy-3-(2-dimethylamino-ethyl)-1*H*-indol-1-yl]phenylmethanone (300 mg, 0.75 mmol) and 4N hydrochloric acid (190 ml, 0.76 mmol) in ethanol (10 ml) was hydrogenated over 10% palladium on carbon (100 mg) at 35 psi for 3 hours. The catalyst was removed by filtration. The filtrate was evaporated, and the residue was partitioned between dichloromethane and saturated aqueous sodium hydrogen carbonate. The aqueous was further extracted with dichloromethane (x2). The combined extracts were dried (K₂CO₃), filtered and evaporated. The residue was purified by chromatography on silica gel, eluting with dichloromethane /methanol/aqueous ammonia (80:8:1) to give the title compound (120 mg, 52%), as a colourless solid; mp 154-156°C. δ_H (360 MHz, CDCl₃) 2.40 (6H, s), 2.70-2.75 (2H, m), 2.83-2.88 (2H, m), 6.86-6.90 (2H, m), 7.04 (1H, s), 7.49-7.61 (3H, m), 7.68-7.71 (2H, m), 8.17 (0.5H, s), 8.20 (0.5H, s). m/z (CI⁺) 309 [MH]⁺. Found: C, 73.22; H, 6.48; N, 8.74. C₁₉H₂₀N₂O₂ . 0.2 H₂O requires C, 73.15; H, 6.59; N, 8.98%.

EXAMPLE 24

20 3-(2-Dimethylamino-ethyl)-5-hydroxy-1*H*-indole-1-carboxylic acid *tert*-butyl ester

a) 5-Benzyloxy-3-(2-dimethylamino-ethyl)-1*H*-indole-1-carboxylic acid *tert*-butyl ester

25 Di-*tert*-butyl dicarbonate (408 mg, 1.9 mmol) was added to a stirred solution of [2-(5-benzyloxy-1*H*-indol-3-yl)ethyl]dimethylamine (500 mg, 1.7 mmol), triethylamine (473 ml, 3.4 mmol) and 4-(dimethylamino)-pyridine (20 mg, 0.2 mmol) in dry dichloromethane (5 ml) at 0°C. The reaction was allowed to warm slowly to ambient temperature. After 3 hours the mixture was poured into saturated aqueous sodium hydrogen carbonate and extracted with dichloromethane (x3). The combined

30

extracts were dried (Na_2SO_4), filtered and evaporated. The residue was purified by chromatography on silica gel, eluting with dichloromethane/methanol/aqueous ammonia (80:8:1) to give the title compound (630 mg, 95%) as a thick oil. δ_{H} (360 MHz, CDCl_3) 1.65 (9H, s), 2.33 (6H, s), 2.58-2.62 (2H, m), 2.79-2.83 (2H, m), 5.12 (2H, s), 6.99 (1H, dd, $J=9.0$ and 2.5 Hz), 7.06 (1H, d, $J=2.5$ Hz), 7.29-7.48 (6H, m), 8.0 (1H, br d).

b) 3-(2-Dimethylamino-ethyl)-5-hydroxy-1H-indole-1-carboxylic acid tert-butyl ester

A solution of 5-benzyloxy-3-(2-dimethylamino-ethyl)-1H-indole-1-carboxylic acid *tert*-butyl ester (303 mg, 0.77 mmol) and 4N hydrochloric acid (192 ml, 0.77 mmol) in ethanol (10 ml) was hydrogenated over 10% palladium on carbon (100 mg) at 40 psi for 4 hours. The catalyst was removed by filtration. The filtrate was evaporated and the residue was partitioned between aqueous sodium hydrogen carbonate and dichloromethane. The aqueous was further extracted with dichloromethane (x2). The combined extracts were dried (K_2CO_3), filtered and evaporated. The residue was purified by chromatography on silica gel, eluting with dichloromethane/methanol/aqueous ammonia (60:8:1) to give the title compound (117 mg, 50%) as a colourless solid. δ_{H} (360 MHz, CDCl_3) 1.64 (9H, s), 2.45 (6H, s), 2.73-2.80 (2H, m), 2.88-2.93 (2H, m), 6.82-6.85 (1H, m), 6.96 (1H, br s), 7.33 (1H, br s), 7.92 (1H, m). m/z (CI^+) 305 $[\text{MH}]^+$. Found: C, 64.07; H, 8.07; N, 8.71. $\text{C}_{17}\text{H}_{24}\text{N}_2\text{O}_3 \cdot 0.8 \text{H}_2\text{O}$ requires C, 64.01; H, 8.15; N, 8.78%.

EXAMPLE 25*N,N*-Dimethyl 2-(1-Benzenesulphonyl-5-benzyloxy-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

5 To a stirred solution of 5-benzyloxytryptamine hydrochloride (2.29 g, 7.56 mmol) in methanol (100 ml) was added sodium methoxide (30% w/v in methanol; 1.44 ml) followed by glacial acetic acid (1.70 ml, 30.2 mmol) and sodium cyanoborohydride (950 mg, 15.12 mmol). A solution of formaldehyde (38% w/v aqueous solution; 1.43 ml) in methanol (25 ml) was
10 then added dropwise over 20 minutes, and the resulting solution was stirred at room temperature for 4 h. 4N Sodium hydroxide (60 ml) was added and the methanol was removed under vacuum. The aqueous residue was extracted with diethyl ether (2 x 250 ml) and the combined organic phases were washed with brine (2 x 50 ml), dried (Na₂SO₄) and
15 concentrated. The residue was dissolved in dichloromethane (120 ml) and 50% aqueous sodium hydroxide (40 ml) was added followed by tetrabutylammonium hydrogen sulphate (258 mg). The mixture was vigorously stirred, benzenesulphonyl chloride (1.45 ml) in dichloromethane (10 ml) was added dropwise over 3 minutes and stirring was continued for
20 30 minutes. Water (120 ml) and diethyl ether (200 ml) were added and the organic phase was decanted off, washed with brine (2 x 50 ml), dried (Na₂SO₄) and concentrated. Flash chromatography of the residue (silica gel, dichloromethane-methanol-ammonia, 95:5:0.5) afforded 2.57 g (78%) of the title compound free base as a colourless thick oil. The oxalate salt
25 was prepared; mp 212-215°C (EtOH); δ_H (360MHz, DMSO-*d*₆) 7.94-7.90 (2H, m), 7.82 (1H, d, *J*=9.0Hz), 7.72-7.64 (2H, m), 7.58 (2H, t, *J*=8.0Hz), 7.48-7.28 (6H, m), 7.06 (1H, dd, *J*=9.0 and 2.4Hz), 5.12 (2H, s), 3.30-3.22 (2H, m), 3.06-2.98 (2H, m), 2.77 (6H, s); *m/z* (ES) 435 (*M*⁺+1). (Found: C, 61.81; H, 5.38; N, 5.48. C₂₅H₂₆N₂O₃S.1.0 C₂H₂O₄ requires: C, 61.82; H, 5.38; N, 5.34%).
30

EXAMPLE 26*N,N*-Dimethyl 2-(1-benzenesulphonyl-5-hydroxy-1*H*-indol-3-yl)ethylamine

A solution of *N,N*-dimethyl 2-(1-benzenesulphonyl-5-benzyloxy-1*H*-indol-3-yl)ethylamine (2.25 g, 4.6 mmol) in absolute ethanol (100 ml) was hydrogenated at 40 psi over 20% palladium hydroxide (600 mg) for 4.5 h. Additional catalyst (1.6 g) was then added and hydrogenation was resumed at 45 psi for 2.5h. The catalyst was removed by filtration, washed with a mixture of dichloromethane-methanol-ammonia (80:20:2; 3 x 70 ml), and the filtrate was concentrated under vacuum. The residue was purified by flash chromatography (silica gel, dichloromethane-methanol-ammonia, 95:5:0.5) to give the title compound as a crystalline solid after trituration with diethyl ether (30 ml); mp 185-188°C; δ_H (360MHz, DMSO- d_6) 9.32 (1H, br s), 7.90-7.84 (2H, m), 7.69 (1H, d, $J=8.7$ Hz), 7.68-7.62 (1H, m), 7.55 (2H, t, $J=8.0$ Hz), 7.48 (1H, s), 6.82 (1H, d, $J=2.2$ Hz), 6.78 (1H, dd, $H=8.8$ and 2.2 Hz), 2.67 (2H, t, $J=7.6$ Hz), 2.47 (2H, t, $J=7.6$ Hz), 2.18 (6H, s); m/z (ES) 345 (M^++1). (Found: C, 62.91; H, 5.86; N, 8.16. $C_{18}H_{20}N_2O_3$ requires: C, 62.77; H, 5.85; N, 8.13%).

EXAMPLE 27*N,N*-Dimethyl 2-(1-benzenesulphonyl-5-cyano-1*H*-indol-3-yl)ethylamineHydrogen Oxalate

To a stirred solution of 3-(2-aminoethyl)-5-cyano-1*H*-indole hydrochloride (Castro, J. L. et al. *J. Med. Chem.* 1994, 37, 3023-3032) (0.1208 g, 0.545 mmol) in methanol (25 ml) was added a 30% w/v solution of sodium methoxide in methanol (0.1 mL, 0.545 mmol), then acetic acid (0.125 ml, 2.18 mmol), followed by sodium cyanoborohydride (68 mg, 1.09 mmol). A 38% w/v aqueous solution of formaldehyde (0.103 ml, 1.31 mmol) in methanol (10 ml) was then added dropwise over 30 min. The mixture was stirred for 3.5 h, then 4N aqueous NaOH (25 ml) was added, and the

methanol was removed in vacuo. The aqueous residue was extracted with diethyl ether (3 x 25 ml), and the combined organic extracts were washed with brine (2 x 20 ml), dried (Na_2SO_4), and evaporated in vacuo to give crude *N,N*-dimethyl 2-(5-cyano-1*H*-indol-3-yl)ethylamine.

5 To this crude product was added dichloromethane (30 ml), 50% aqueous NaOH (2 ml), and tetrabutylammonium hydrogen sulphate (19 mg, 0.0545 mmol). Benzenesulphonyl chloride (0.104 ml, 0.817 mmol) was then added dropwise over 3 min whilst stirring magnetically, and the mixture was stirred for 30 min. Ethyl acetate (50 ml) and water (25 ml)
10 was then added, and the organic layer was washed with brine (2 x 25 ml), dried (Na_2SO_4), and evaporated in vacuo. The residue was purified by flash chromatography (silica gel, CH_2Cl_2 -MeOH- $\text{NH}_3(\text{aq})$; 95:5:0.5) to yield 0.132 g (66%) of the title compound free base. The oxalate salt was prepared in EtOH-Et₂O as a white solid; mp 213-220°C; δ_{H} (360 MHz, *d*₆-DMSO) 2.78
15 (6H, s), 3.10 (2H, m), 3.31 (2H, m), 7.63 (2H, t, *J*=7.5 Hz), 7.74 (1H, t, *J*=7.5 Hz), 7.78 (1H, dd, *J*=8.6, 1.4 Hz), 7.98 (1H, s), 8.03 (2H, d, *J*=7.4 Hz), 8.11 (1H, d, *J*=8.4 Hz), 8.31 (1H, s); *m/z* (ES^+) 354 (MH^+). Found: C, 57.01; H, 4.83; N, 9.22. $\text{C}_{19}\text{H}_{19}\text{N}_3\text{O}_2\text{S} \cdot \text{C}_2\text{H}_2\text{O}_4$ requires C, 56.88; H, 4.77; N, 9.48%.

20

EXAMPLE 28

N,N-Dimethyl 2-(1-benzenesulphonyl-5-methoxy-2,3-dihydro-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

25 a) *N,N*-Dimethyl 2-(5-methoxy-2,3-dihydro-1*H*-indol-3-yl)ethylamine

A mixture of 5-methoxy-*N,N*-dimethyltryptamine (0.200g, 0.939 mmol) and palladium on carbon (10%, 200 mg) in 5 M aqueous HCl (50 ml) was hydrogenated at 50 psi overnight. The catalyst was removed by filtration, and the filtrate was extracted with dichloromethane (3 x 25 ml).
30 The combined organic extracts were evaporated in vacuo and the residue was purified by flash chromatography (silica gel, CH_2Cl_2 -MeOH- $\text{NH}_3(\text{aq})$;

92:8:0.8) to give 94 mg (47%) of the title compound; δ_{H} (250 MHz, CDCl_3) 1.71 (1H, m), 1.99 (1H, m), 2.26 (6H, s), 2.37 (2H, m), 3.20 (1H, t, $J=8.0$ Hz), 3.27 (1H, m), 3.68 (1H, t, $J=8.0$ Hz), 3.75 (3H, s), 6.60-6.61 (2H, m), 6.73 (1H, m).

5

b) *N,N*-Dimethyl 2-(1-benzenesulphonyl-5-methoxy-2,3-dihydro-1*H*-indol-3-yl)ethylamine Hydrogen Oxalate

Following a similar procedure to that described in Example 6, but using *N,N*-dimethyl 2-(5-methoxy-2,3-dihydro-1*H*-indol-3-yl)ethylamine
10 instead of 5-methoxy-*N,N*-dimethyltryptamine, the title compound was prepared in 57% yield as a white solid; mp 183-185°C; δ_{H} (360 MHz, d_6 -DMSO) 1.47 (1H, m), 1.94 (1H, m), 2.68 (6H, s), 2.96 (2H, m), 3.14 (1H, m), 3.63 (1H, m), 3.71 (3H, s), 4.06 (1H, m), 6.82 (2H, m), 7.42 (1H, d, $J=9.4$ Hz), 7.57 (2H, t, $J=7.3$ Hz), 7.68 (1H, t), 7.78 (2H, d, $J=7.6$ Hz); m/z (ES^+)
15 361 (MH^+). Found: C, 54.29; H, 5.56; N, 5.97. $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_3\text{S} \cdot \text{C}_2\text{H}_2\text{O}_4 \cdot 0.7\text{H}_2\text{O}$ requires C, 54.46; H, 5.96; N, 6.05%.

EXAMPLE 29

20 *trans*-4-Dimethylamino-1-(4-methylbenzenesulphonyl)-1,3,4,5-tetrahydro-benz[*cd*]indol-5-ol. Hydrogen Oxalate

A suspension of (\pm)-4-azido-3,4-dihydro-1-[(methylphenyl)sulphonyl]benz[*cd*]indol-5(1*H*)one (*J. Chem. Soc. Perkin Trans. I*, 1973, 438-442) (11.9 g) in absolute ethanol (400 ml) and
25 concentrated hydrochloric acid (11.9 ml) was hydrogenated at 25 psi over 10% palladium on carbon (1.2 g) for 6 h. The mixture was diluted with absolute ethanol (300 ml), 10% palladium on carbon (2 g) was added and hydrogenation was resumed for another 16 h at 25 psi. The catalyst was removed by filtration, washed with absolute ethanol (3 x 125 ml) and the
30 filtrate was concentrated under vacuum. The residue was crystallised from ethanol to give 4.55 g of (\pm)-4-amino-3,4-dihydro-1-

[(methylphenyl)sulphonylbenz[cd]indol-5(1H)one hydrochloride as a white solid.

To a solution of sodium borohydride (1 g) in absolute ethanol (75 ml) was added dropwise *via* cannula a suspension of the above aminoketone hydrochloride (4.55 g) in absolute ethanol (150 ml) over 15 minutes, keeping the internal temperature below 3°C. The resulting mixture was stirred at -2°C for 1h 40 min under nitrogen before it was carefully acidified with 2N hydrochloric acid (28 ml). The ethanol was removed under vacuum and the residue was basified with 2N sodium hydroxide (35 ml). The products were extracted with ethyl acetate (3 x 150 ml) and the combined organic phases were washed with brine (50 ml), dried (Na₂SO₄) and concentrated. Flash chromatography of the residue (silica gel, dichloromethane-ethanol-ammonia, 170:28:3) gave 3.99 g (97%) of 4-amino-1-(4-methyl)benzenesulphonyl-1,3,4,5-tetrahydrobenz[cd]indol-5-ol as a 90:10 mixture of trans and cis isomers; *m/z* 342 (M⁺).

A solution of the above aminoalcohol (3.97 g, 11.59 mmol) in absolute ethanol (150 ml), aqueous formaldehyde (38% w/v; 9.2 ml) and glacial acetic acid (92 ml) was hydrogenated at 30 psi over 10% palladium on carbon (1.56 g) for 65 h. The catalyst was filtered off, washed with absolute ethanol and the filtrate was concentrated under vacuum. The residue was dissolved in dichloromethane (40 ml) and washed with 10% aqueous sodium hydrogen carbonate (100 ml). The aqueous layer was extracted with dichloromethane (2 x 120 ml) and the combined organic phases were dried (Na₂SO₄) and concentrated. Flash chromatography of the residue (silica gel, dichloromethane-ethanol-ammonia, 85:15:0 to 1750:225:25) gave 3.68 g (86%) of the title compound free base together with 400 mg of the corresponding *cis*-isomer. The oxalate salt was prepared from ethanol-diethyl ether; mp 123-126°C; δ_H (360MHz, DMSO-*d*₆) 7.86 (2H, d, *J*=8.4Hz), 7.76 (1H, d, *J*=8.1Hz), 7.55 (1H, s), 7.44-7.36 (3H, m), 7.30 (1H, d, *J*=7.4Hz), 5.14 (1H, d, *J*= 9.3Hz), 3.54-3.38 (1H, m), 3.26-3.16 (1H, m), 3.12-2.96 (1H, m), 2.78 (6H, s), 2.32 (3H, s); *m/z* (EI) 371

(M⁺+1). (Found: C, 57.16; H, 5.36; N, 5.99. C₂₀H₂₂N₂O₃S.1.0 C₂H₂O₄ requires: C, 57.38; H, 5.25; N, 6.08%). [*J Chem Soc. Perkin Trans I*, 1973, 438-442]

5

EXAMPLE 30

4-Dimethylamino-5-methoxy-1-(4-methylbenzenesulphonyl)-1,3,4,5-tetrahydro-benz[cd]indole Hydrogen Oxalate

To a solution of the product from Example 29 (free base; 700 mg) in
10 anhydrous tetrahydrofuran (10 ml) and anhydrous dimethylformamide (3 ml) was added sodium hydride (60% dispersion in oil; 91 mg) followed, after 10 minutes, by benzyltrimethylammonium chloride (35 mg) and methyl 3-bromopropionate. After stirring at room temperature for 2.5 h, additional sodium hydride was added and stirring was continued for 16 h.
15 Water (50 ml) was added and products were extracted with dichloromethane (3 x 50 ml), then dried (MgSO₄) and concentrated. Flash chromatography of the residue (silica gel, dichloromethane-ethanol, 95:5) gave 30 mg of the title compound free base. The oxalate salt was prepared and crystallised from isopropanol; mp 133-136°C; δ_H (360MHz, DMSO-d₆)
20 7.86 (2H, d, J=8.4Hz), 7.80 (1H, d, J=8.1Hz), 7.52 (1H, s), 7.42-7.34 (3H, m), 7.29 (1H, d, J=7.3Hz), 4.89 (1H, d, J=6.5Hz), 3.56-3.48 (4H, m and s), 3.08 (2H, d, J=5.6Hz), 2.52 (6H, s), 2.31 (3H, s); m/z 384 (M⁺). (Found: C, 58.01; H, 5.84; N, 5.77. C₂₁H₂₄N₂O₃S.1.0 C₂H₂O₄ requires: C, 58.22; H, 5.52; N, 5.90%).

25